



SOUTH COAST
UROGYNECOLOGY

THE LAGUNA LASER CENTER

Red M. Alinsod, M.D.,
FACOG, FACS, ACGE
red@urogyn.org

Maggie Carpio, P.A.-C
maggie@urogyn.org

Toll free (877) 4-UROGYN
Main (949) 499-5311
Fax (949) 499-5312

Consent for LUMIGAN

LUMIGAN contains:

Active: bimatoprost 0.3 mg/lm

Preservative: Benzalkonium chloride 0.05mg/ml

Inactives: Sodium chloride, sodium phosphate, dibasic, citric acid and purified water.

Indication and Usage

LUMIGAN (bimatoprost ophthalmic solution) 0.03% is indicated for the reduction of elevated intraocular pressure in patients with open angle glaucoma or ocular hypertension.

Contraindications

LUMIGAN is contraindicated in patients with hypersensitivity to bimatoprost or any other ingredient in this product.

Warnings

LUMIGAN has been reported to cause changes to pigmented tissues. The most frequently reported changes have been increased pigmentation of the iris, periorbital tissue (eyelid) and eyelashes, and growth of eyelashes. Pigmentation is expected to increase as long as LUMIGAN is administered. After discontinuation of LUMIGAN pigmentation of the iris is likely to be permanent while pigmentation of the periorbital tissue and eyelash changes have been reported to be reversible in some patients. Patients who receive treatment should be informed of the possibility in increased pigmentation. The effects of increased pigmentation beyond 5 years are not known.

Precautions

LUMIGAN may gradually change eyelashes and vellus hair in the treated eye; these changes include increased length, thickness and number of lashes. Eyelash changes are usually reversible upon discontinuation of treatment.

LUMIGAN should be used with caution in patients with active intraocular inflammation (e.g., uveitis).

Macular edema, including cystoid macular edema, has been reported during treatment with bimatoprost ophthalmic solution. LUMIGAN should be used with caution in aphakic patients, in pseudophakic patients with a torn posterior lens capsule, or in patients with known risk factors for macular edema.

Contact lenses should be removed prior to instillation of LUMIGAN and may be reinserted 15 minutes following its administration.

If you should develop any ocular reactions, particularly conjunctivitis and eyelid reactions, please discontinue use and contact Maggie Carpio P.A.-C at The Laguna Laser Center at 949-499-5311.

I have been advised of the indications for use of LUMIGAN and possible risks and side effects.

I have been advised of the indications for use of LUMIGAN and possible risks and side effects. I am not aware that I am pregnant. I have discussed all my concerns and my questions have been answered.

Patient Signature _____ **Date** _____

Witness Signature _____ **Date** _____

LUMIGAN initial 2-1/2cc lasts about 3 months
Instructions

You will need an eyeliner brush that can be purchased at a beauty supply store.

- Wash hands and try to keep as sterile as possible.
- Remove dropper cap, place product on your hand, dip the brush in product and apply with eyeliner brush as you would eyeliner. (Placing the product on your hand keeps the rest of your solution clean.)
- Apply once every night for 10-12 weeks. Results are usually seen after 4 weeks but may take up to 2-3 months. Results may vary for each person.
- After the initial 10-12 weeks, maintain growth by applying once a week.
- Rinse your brush after each application.
- If irritation occurs, discontinue use until the irritation clears, then start again. If you continue to experience irritation or allergic reaction, discontinue use altogether, and consult your doctor.

